

Don't forget subterranean ecosystems in climate change agendas

To the Editor — Climate change is a mounting threat to biodiversity, affecting multiple ecosystem services critical for human wellbeing, and hence the aim of most Sustainable Development Goals¹. To address this herculean challenge, several global initiatives at the crossroads of science and policy have been advanced to implement

bold and effective post-2020 biodiversity and climate change agendas². For example, the *Half-Earth Project* (www.half-earthproject.org) or the *Global Safety Net*³ target 50% of Earth to be formally protected to halt biodiversity loss, reduce CO₂ emissions and enhance natural carbon sequestration. To be optimally effective, these programmes

should aim to include the maximum breadth of species, habitats and ecosystem services. Upon evaluating the diverse post-2020 conservation agendas, however, we were struck by a curious omission. Subterranean ecosystems — likely the most widespread non-marine environment on Earth^{4,5} — were unfortunately overlooked.




Fig. 1 | Biodiversity, ecosystem services and current status of protection of subterranean ecosystems. The extent of subterranean ecosystems was calculated by combining geospatial layers of the primary strata supporting subterranean habitats — karst and lava fields. An estimation of subterranean diversity and examples of ecosystem services were based on recent literature^{18–21}.

Whereas we do not know the true global extent of the subterranean biome⁵, a rough estimation using geospatial data of relevant geological substrates suggests that it covers at least 19% of the terrestrial surface⁶. Of this, just 6.9% (Fig. 1) overlaps with currently protected areas⁷, which is a rather low percentage for an ecosystem type. Importantly, this estimate does not account for the vertical dimension of subterranean ecosystems.

The omission of subterranean ecosystems from global biodiversity and climate change agendas marginalizes their ecological and economic importance. From a biodiversity point of view, these systems harbour a great diversity of specialized organisms that are of interest from both a conservation and evolutionary perspective (Fig. 1). Subterranean species are often short-range endemics (occurring within a single cave or geological formation), and some represent ancestors of faunas that disappeared from surface habitats. Thus, they account for a considerable proportion of global taxonomic, phylogenetic and functional diversity that is currently imperilled by human activities⁴, especially climate change⁸.

Being adapted to environments with limited seasonality, stable temperatures and a near-water-saturated atmosphere with a negligible evaporation rate⁸, most of these species have lost the physiological mechanisms to withstand rapid environmental fluctuations^{9,10}. Although a lag effect may be expected with subterranean temperatures compared to adjacent surface ecosystems, climate change is predicted to alter conditions that these sensitive species currently rely upon⁸. Temperature increase, drought-related stressors and ecological shifts of surface ecosystems (for example, desertification) are all projected to alter hydrological regimes and pervert subterranean conditions — rendering some suitable habitats unsuitable. When considering that most specialized subterranean species are unable to escape the effects of climate change due to geographical, phenological and behavioural constraints, subterranean ecosystems may represent dead-end traps for their obligate inhabitants⁸.

By threatening subterranean biodiversity, climate change will in turn affect the provisioning of crucial ecosystem services to humankind⁴ (for examples of services, see Fig. 1). Perhaps most importantly, an estimated 95% of the world's available liquid freshwater supply (that is, excluding ice

caps) is groundwater, and over one-quarter of the human population is completely or partially dependent on it¹¹. Bacteria and invertebrates play a key role in maintaining clean groundwaters as they are responsible for the carbon turnover and the attenuation and degradation of harmful contaminants, and may even eliminate pathogenic viruses and microorganisms¹². Globally, the human demand for groundwater is ~3.5 times the actual volume of aquifers¹³, and this situation will only worsen as the human population keeps growing, and the intensification of drought events will increase the demand of groundwaters for agricultural irrigation^{11,14}. The combination of these and other climate-change-related stressors, such as saltwater intrusion in coastal aquifers, are predicted to eliminate or constrict aquatic subterranean freshwater habitats, threatening subterranean biodiversity and reducing water availability for human use^{14,15}.

Another well-documented ecological service is provided by cave-roosting bats. Nearly 150 bat species use subterranean habitats (IUCN Red List, version 2020-3). Of these, many provide important ecosystem services, including seed dispersal and pollination of both timber and food crops as well as the consumption of agricultural pests¹⁶. Rising temperatures may cause current summer roost caves to become too warm, impeding maternity activities, as well as arousing bats from hibernation during times of low prey availability¹⁷. These changes could facilitate further population declines, which would spur a reduction in bat ecosystem services^{18–21}.

As subterranean ecosystems have been lamentably overlooked, perhaps the aphorism of 'out of sight, out of mind' has driven both scientific interest and policy decisions. It is therefore critical to develop a strategic plan to monitor subterranean habitats and biodiversity, which can be integrated into current global conservation targets. This strategy should include the identification of a suite of bioclimatic indicators to safeguard key ecosystem services and highly biodiverse subterranean communities.

This year marks the *International Year of Caves and Karst* (<http://iyck2021.org>). Serendipitously, we hope 2021 will also represent an inflection point in global decision-making concerning subterranean habitats. It is high time these critically important ecosystems assume their rightful place in global biodiversity

conservation strategies and climate change mitigation agendas. □

David Sánchez-Fernández^{1,6},
Diana M. P. Galassi², J. Judson Wynne³,
Pedro Cardoso⁴ and
Stefano Mammola^{1,4,5,6}✉

¹Departament of Ecology and Hydrology, University of Murcia, Murcia, Spain. ²Department of Life, Health & Environmental Sciences, University of L'Aquila, L'Aquila, Italy. ³Department of Biological Sciences, Center for Adaptable Western Landscapes, Northern Arizona University, Flagstaff, AZ, USA. ⁴LIBRe—Laboratory for Integrative Biodiversity Research, Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland. ⁵DarkMEG—Molecular Ecology Group, Water Research Institute (IRSA), National Research Council of Italy (CNR), Verbania Pallanza, Italy. ⁶These authors contributed equally: David Sánchez-Fernández, Stefano Mammola.

✉e-mail: stefano.mammola@helsinki.fi

Published online: 20 May 2021
<https://doi.org/10.1038/s41558-021-01057-y>

References

1. *Nat. Clim. Change* **9**, 491 (2019).
2. Arnet, A. et al. *Proc. Natl Acad. Sci. USA* **117**, 30882–30891 (2020).
3. Dinerstein, E. et al. *Sci. Adv.* **6**, eabb2824 (2020).
4. Mammola, S. et al. *BioScience* **69**, 641–650 (2019).
5. Ficetola, G. F., Canedoli, C. & Stoch, F. *Conserv. Biol.* **33**, 214–216 (2019).
6. Chen, Z. et al. *World Karst Aquifer Map (WHYMAP WOKAM)* (BGR, IAH, KIT & UNESCO, 2017); https://doi.org/10.25928/b2.21_sfkq-r406
7. *World Database on Protected Areas* (IUCN & UNEP-WCMC, 2016).
8. Mammola, S. et al. *Anthr. Rev.* **6**, 98–116 (2019).
9. Pallarés, S. et al. *Anim. Conserv.* <https://doi.org/10.1111/acv.12654> (2021).
10. Castaño-Sánchez, A., Hose, G. C. & Reboreira, A. S. *P. Sci. Rep.* **10**, 12328 (2020).
11. Taylor, R. G. et al. *Nat. Clim. Change* **3**, 322–329 (2012).
12. Griebler, C. & Avramov, M. *Freshw. Sci.* **34**, 355–367 (2015).
13. Gleeson, T., Wada, Y., Bierkens, M. F. P. & van Beek, L. P. H. *Nature* **488**, 197–200 (2012).
14. Wu, W. Y. et al. *Nat. Commun.* **11**, 3710 (2020).
15. Famiglietti, J. S. *Nat. Clim. Change* **4**, 945–948 (2014).
16. Frick, W. F. et al. *Ann. N. Y. Acad. Sci.* **1469**, 5–25 (2020).
17. Browning, E. et al. *Mamm. Rev.* <https://doi.org/10.1111/mam.12239> (2021).
18. Deharveng, L. & Bedos, A. In *Cave Ecology* 107–172 (Springer, 2019).
19. Stein, H. et al. *Sci. Rep.* **2**, 673 (2012).
20. Culver, D. C. & Holsinger, J. R. *Nat. Speleol. Soc. Bull.* **54**, 79–80 (1992).
21. Magnabosco, C. et al. *Nat. Geosci.* **11**, 707–717 (2018).

Acknowledgements

This study is supported by the project CAWEB (Testing macroecological theory using simplified systems — 882221), funded by Marie Skłodowska-Curie Individual Fellowships (H2020-MSCA-IF-2019) to S.M. D.S.-F. is funded by the Ramón y Cajal programme (RYC2019-027446-I) from the Spanish Ministry of Science and Innovation. DMPG is funded by the LIFE Programme, AQUALIFE LIFE12 BIO/IT/000231.

Competing interests

The authors declare no competing interests.